

Increased Frequency of Ig Heavy-Chain HS1,2-A Enhancer *2 Allele in Dermatitis Herpetiformis, Plaque Psoriasis, and Psoriatic Arthritis

Rossella Cianci^{1,7}, Vincenzo Giambra^{2,7}, Claudia Mattioli², Maria Esposito³, Giovanni Cammarota¹, Graziella Scibilia⁴, Giuseppe Magazzù⁴, Ambrogio Orlando⁵, Giancarlo Sandri⁶, Luca Bianchi³, Giovanni Battista Gasbarrini¹, Franco Pandolfi¹ and Domenico Frezza²

The enhancer DNase-hypersensitive region 1,2 (HS1,2), a member of the Ig heavy-chain 3' regulatory region (3'RR) cluster, is active in human B cells transfected with reporter genes and in mouse is activated in late maturation. HS1,2-A contains binding sites for several transcription factors. There are four known alleles, that is, *1, *2, *3, and *4, which differ in their lengths in transcription factor binding. We showed that in celiac disease the frequency of the *2 allele is increased. Both dermatitis herpetiformis (DH) and psoriasis can be associated with different frequencies with celiac disease. Thus, we further investigate the frequency of allele *2 in DH, plaque psoriatic, and psoriatic arthritis patients. HS1,2-A allele frequencies were investigated in 37 DH, 61 plaque psoriatic, 28 psoriatic arthritis patients, and 265 healthy donors, age- and sex-matched, from the same geographical area. The frequency of the *2 allele changes from 0.39 in controls to 0.63 in DH, 0.59 in plaque psoriasis and 0.75 in psoriatic arthritis (P between 10^{-4} – 10^{-5}). Our data evidence an increased frequency of the *2 allele of HS1,2-A in these cutaneous immune-related disorders. We suggest a related genetic predisposition in these pathogeneses.

Journal of Investigative Dermatology (2008) **128**, 1920–1924; doi:10.1038/jid.2008.40; published online 6 March 2008

INTRODUCTION

Susceptibility to immune diseases is considered a multigenic phenotype affected by a variety of genetic and environmental or stochastic factors (Shepshelovich and Shoenfeld, 2006). The abnormal immune response can be triggered by interactions between susceptibility genes and environmental factors. The most potent genetic influence on susceptibility to immune-disorders is the major histocompatibility complex. However, there also are non-major histocompatibility complex susceptibility alleles related to immune diseases. Various epistatic interactions are predicted to involve the multiple

genes required for disease development (Wandstrat and Wakeland, 2001).

Accordingly, we predict that certain immunological pathways are common to multiple immune diseases, and other pathophysiological mechanisms are specific to a particular disease (Wandstrat and Wakeland, 2001). For example, the presence of a specific antibody in an individual's serum combined with disease-prone major histocompatibility complex haplotypes or other susceptibility loci exponentially increases the risk for that person to develop an immune disease in the future (Shepshelovich and Shoenfeld, 2006).

Plaque psoriasis (OMIM 177900) and psoriatic arthritis (OMIM 607507), together with dermatitis herpetiformis (DH) (OMIM 601230), are multi-factorial cutaneous diseases with familial incidence and immune dependence (Karell *et al.*, 2002; Bowcock, 2005). Celiac disease and DH are closely related diseases, which share a genetic background and a common pathogenesis characterized by gluten sensitivity and production of autoantibodies (Hervonen *et al.*, 2000; Karell *et al.*, 2002). Patients with DH, which is sometime referred to as the cutaneous phenotype of celiac disease, have asymptomatic or frank celiac disease and a skin rash with anti-transglutaminase (TGM) cutaneous IgA deposits. Celiac disease and DH patients both show anti-TGM type 2 autoantibodies. Sardy *et al.* (2002) have shown a difference in the pattern of anti-TGM antibodies. In fact, in addition to anti-TGM type 2 antibodies, DH patients develop a further

¹Institute of Internal Medicine, Catholic University, Rome, Italy; ²Department of Biology "Enrico Caffè", Tor Vergata University, Rome, Italy; ³Department of Dermatology, Tor Vergata University Hospital, Rome, Italy; ⁴Department of Paediatrics, University Hospital, Messina, Italy; ⁵Department of Internal Medicine, "V Cervello" Hospital, Palermo, Italy and ⁶Nutrition Unit, S Eugenio Hospital, Rome, Italy

⁷These authors contributed equally to this work

The work was done in Rome and the samples were collected from different Clinics of Roma, Messina, and Palermo, Italy

Correspondence: Professor Domenico Frezza, Department of Biology "Enrico Caffè", Tor Vergata University, Viale dell'Università, Rome 00133, Italy. E-mail: frezza@uniroma2.it

Abbreviations: DH, dermatitis herpetiformis; HS, DNase-hypersensitive region; HS1,2-A, DNase-hypersensitive A region 1,2; 3'RR, 3' Ig heavy-chain regulatory region; TGM, transglutaminase

Received 8 March 2007; revised 6 November 2007; accepted 14 December 2007; published online 6 March 2008

class of high-affinity antibodies against epidermal TGM or TGM type 3. Therefore, TGM3 is the dominant autoantigen in DH, explaining why skin symptoms rather than intestinal symptoms appear in a proportion of patients with gluten-sensitive disease (Sardy *et al.*, 2002).

Plaque psoriasis, a clinically heterogeneous disease with inherited susceptibility, is considered a T-cell-mediated inflammatory skin disease, but humoral immune processes have also been evaluated. Plaque psoriasis has been variously reported to be associated with IgA-related disorders such as celiac disease, dermatitis herpetiformis, linear IgA sub-epidermal bullous dermatosis, IgA nephropathy, or IgA necrotic renal glomerular vasculitis (Kallel Sellami *et al.*, 2006; Lapeyre *et al.*, 2006; Zadrazil *et al.*, 2006). Up to 16% of psoriatic patients show the presence of IgG and IgA anti-endomysial, anti-gliadin, and anti-TGM antibodies (Woo *et al.*, 2004). These celiac disease-associated antibodies in psoriatic patients are apparently correlated with greater disease activity. In particular, IgA anti-tissue TGM antibodies seems to represent a lesion-associated event (Borg, 1997). A good correlation between serum and synovial fluid anti-tissue TGM levels was found in all inflammatory bowel patients and in arthropathic lesions (Caspi *et al.*, 2006). Furthermore, psoriatic arthritis patients seem to have an increased number of lymphocytes in the duodenal mucosa and a highly significant increase in intraepithelial lymphocytes in comparison with plaque psoriasis patients (Lindqvist *et al.*, 2006). Taken together, these data strongly support the idea of IgA deregulation in these skin diseases and more generally, an immune-alteration.

The 3' Ig heavy-chain regulatory region (3'RR) enhancer complex (Pettersson *et al.*, 1990; Chen and Birshstein, 1997; Mills *et al.*, 1997) is active in murine class switch

recombination and B-cell plasma cells (Andersson *et al.*, 2000; Pinaud *et al.*, 2001). The 3'RR is present through many species from mouse to apes and the orthologous enhancers conserved (Harindranath *et al.*, 1998; Stavnezer and Amemiya, 2004; Sepulveda *et al.*, 2005).

In humans, the 3'RR is duplicated; a copy lies downstream of each of constant α -1 and constant α -2 genes (Chen and Birshstein, 1997; Mills *et al.*, 1997; Pinaud *et al.*, 1997). The human 3'RRs contain three enhancers: DNase-hypersensitive region 3 (HS3), HS1,2, and HS4. The central HS1,2 enhancers are polymorphic (see Figure 1; Mills *et al.*, 1997; Giambra *et al.*, 2005). A significant correlation of the *2 allele of DNase-hypersensitive A region 1,2 (HS1,2-A), downstream of α -1 gene, was shown in patients with IgA nephropathy, and celiac disease and systemic sclerosis (Aupetit *et al.*, 2000; Frezza *et al.*, 2004, 2007).

Therefore, we compared the frequencies of four alleles of the HS1,2-A enhancer in patients with DH, plaque psoriasis, and psoriatic arthritis with a healthy sample of the same geographical area. A significant correlation was observed in all three disorders, with an increase in *2 allele frequency.

RESULTS

The data reported in Table 1 show the frequencies of the four alleles of the enhancer HS1,2-A in the group of patients with dermatitis herpetiformis, plaque psoriasis, psoriatic arthritis, and in a control group selected from the same geographical area and not related at least for three generations. The distribution of the four allele's frequencies is similar in these three pathological groups, but different with respect to the control sample. In the control and pathological samples, the observed genotypes were in agreement with Hardy-Weinberg expected values as reported in Table 1.




Figure 1. The human constant genes heavy-chain locus. (a) The duplication of four constant genes and relative regulatory region 3'RR-1 and 3'RR-2. (b) Enlarged view of the structure of HS1,2 alleles *1 and *2 with the repeated elements of 38 bp and spacers of 17 bp; the triangle shows the position of the consensus of NF- κ B and SP1, and the parallelogram shows the consensus for OCT1. (c) Alignment of allele *1 and *2 sequences and the difference due to the duplication of the 38-mer and internal spacer. The consensus for OCT1, NF- κ B, and SP1 transcription factors are highlighted by square frames.

Table 1. Frequency \pm SE of HS1,2-A alleles in patients with DH, plaque psoriasis, psoriatic arthritis, and healthy control, and Hardy–Weinberg evaluation

Alleles	Dermatitis herpetiformis	Plaque psoriasis	Psoriatic arthritis	Control
*1A	0.297 \pm 0.053 (22)	0.344 \pm 0.043 (42)	0.214 \pm 0.054 (12)	0.449 \pm 0.022 (238)
*2A	0.635 \pm 0.070 (47)	0.590 \pm 0.044 (72)	0.750 \pm 0.057 (42)	0.392 \pm 0.021 (208)
*3A	0.014 \pm 0.014 (1)	0.008 \pm 0.008 (1)	0.018 \pm 0.017 (1)	0.059 \pm 0.010 (31)
*4A	0.054 \pm 0.026 (4)	0.058 \pm 0.021 (7)	0.018 \pm 0.017 (1)	0.100 \pm 0.013 (53)
Genomes	74	122	56	530
Degree of freedom	1	1	1	1
χ^2	4.999	2.088	0.326	1.685
P	0.05 < P < 0.025	0.50 < P < 0.25	0.75 < P < 0.50	0.25 < P < 0.10

DH, dermatitis herpetiformis; HS1,2-A, DNase-hypersensitive A region 1,2.
The number of observed genomes is indicated in parentheses.

Table 2. Test of significance of HS1,2-A allelic frequency differences among DH, plaque psoriasis, or psoriatic arthritis patients compared with the control

	Dermatitis herpetiformis	Plaque psoriasis	Psoriatic arthritis
Degree of freedom	3	3	3
χ^2	16.46	18.63	22.56
P	0.0008	0.0002	< 10^{-5}

DH, dermatitis herpetiformis; HS1,2-A, DNase-hypersensitive A region 1,2.

The comparisons of the allelic frequencies between pathological and healthy groups measured by a χ^2 significance test showed very low *P*-values for DH, plaque psoriasis, and psoriatic arthritis ($P < 8 \times 10^{-4}$, $P < 2 \times 10^{-4}$, and $P < 10^{-5}$, respectively), emphasizing the strong divergence from healthy group (see Table 2). In these three diseases, the frequencies of *1, *3, and *4 alleles are decreased compared with the control. Moreover, there is a dramatic increase in *2 allele frequency, which is 0.63, 0.59, 0.75 in DH, plaque psoriasis, and psoriatic arthritis, respectively, and 0.39 in the control, with an increase of 1.62, 1.50, and 1.91-fold, respectively.

These results indicate a correlation between the *2 allele's frequency increase and DH, plaque psoriasis, and psoriatic arthritis. Therefore, we measured the odds ratios with the corresponding confidence intervals, as reported in Table 3. A similar trend characterizes the three diseases and suggests that the presence of *2 allele associates to a clinical risk in developing the immune-disorders (odds ratio: 2.1, 2.2, and 3.6, respectively).

DISCUSSION

In this report, we provide evidence that the HS1,2-A enhancer *2 allele's frequency is increased in DH, plaque psoriasis, and arthritic psoriasis. These data suggest a possible role for a particular allele of the enhancer HS1,2-A in these

Table 3. Risk to develop DH, plaque psoriasis, or psoriatic arthritis associated with the frequency of the *2A allele

Dermatitis herpetiformis versus control
Degree of freedom=1 $\chi^2=8.291^1$ <i>P</i> =0.004 OR=2.136 CI=1.29–3.53 ²
Plaque psoriasis versus control
Degree of freedom=1 $\chi^2=15.025^1$ <i>P</i> =0.0001 OR=2.229 CI=1.50–3.33 ²
Psoriatic arthritis versus control
Degree of freedom=1 $\chi^2=17.198^1$ <i>P</i> <0.0001 OR=3.681 CI=1.96–6.90 ²

CI, confidence interval; DH, dermatitis herpetiformis; OR, odds ratio.
 χ^2 -Test and OR with CI of the different groups of patients versus control subjects for the *2A allele.

¹With Yates correction.

²Using the approximation of Woolf.

skin disorders. However, it should be noted that the genetic predisposition might not directly relate to cutaneous manifestations. The higher allele *2 frequency in psoriatic patients with arthritis might rather suggest a role for this allele also in the arthritic manifestations. We predict that the increased frequency of *2 allele is not a specific marker of these three pathological conditions, but a marker of a differential response of Ig induction. The *2 allele could function as a magnifier of Ig production that, in a physiological condition, could give a favorable response. In fact in the European population, this allele is present at such a high percent as 39%, impossible if there were not advantages to its presence at least in heterozygotes (Giambra *et al.*, 2006). The frequencies of *2 allele in the three pathologies compared with the control are different, with a highly significant χ^2 -value. The frequency is higher in DH and psoriatic arthritis than in plaque psoriasis; DH and psoriatic arthritis are in fact more rarely observed than plaque psoriasis. This difference could be attributed to the fact that a restricted number of factors are necessary and sufficient to trigger the onset of DH and psoriatic arthritis compared with those of plaque psoriasis, where many independent factors can be sufficient, and not all are necessary, to turn on the disease.

As far as the role of HS1,2-A is concerned, *in vitro* studies demonstrated an increased enhancer activity on the luciferase gene by the HS1,2-A allele *2 compared with *1 as well as for all the alleles harboring a higher number of repeats in respect to the shorter ones (Denizot *et al.*, 2001). The role of HS1,2-A allele *2 may be then important in modifying immune or inflammatory responses in our patients. The most evident difference of allele *2 in regards to allele *1 is the presence of an extra consensus site for NF- κ B and SP1 (see Figure 1). On the basis of the immunological alterations present in the disorders, the *2 allele prompts Ig production to a stronger response either in terms of activation of class switch or germinal transcription. These results may identify a role for genetic polymorphism in the immune regulation and its influence in B-cell activity. The analysis of the haplotypes of the entire 3'-regulatory region will provide more information on the role of this *cis*-acting region in immune-diseases.

In conclusion, our data add DH, plaque psoriasis, and psoriatic arthritis to the growing list of disorders associated with Ig deregulation, where the enhancer HS1,2-A may be involved. Further studies are needed to understand its role in the pathogenesis of the diseases and their progression. It is interesting to note that as of now none of the whole-genome studies in psoriasis have picked up this association. A replication of the findings by independent research groups with larger sample sizes will be necessary to confirm these data.

MATERIALS AND METHODS

Patients

Thirty-seven DH patients were studied (13 males, 24 females, mean age 36.87 ± 17.68 years). Diagnosis was based on standard clinical and laboratory criteria. Granular IgA deposits in the dermal papillae of perilesional skin were evaluated. Celiac disease was associated in 28/37 (8 males, 20 females, mean age 35.50 ± 12.65 years). We have also enrolled 61 patients with plaque psoriasis (39 males, 22 females, mean age 43.91 ± 16.71 years) and 28 with psoriatic arthritis (16 females, 12 males, mean age 47.06 ± 13.70 years). Diagnosis was based on clinical criteria. To exclude association with celiac disease, all psoriatic patients were put on blood tests to assess the presence of anti-endomysial antibody (anti-EMA) antibodies. Two patients with anti-EMA antibodies were excluded.

The control group consists of 265 healthy Italian blood donors, age- and sex-matched, from the same geographical areas. Genomic DNA was extracted from total blood according to standard procedures. All participants gave written informed consent. The study was conducted according to the Declaration of Helsinki Principles and the medical ethical committees of Clinics of Roma, Messina, and Palermo approved the study.

PCR assay

To estimate the frequencies of the four alleles of HS1,2-A (GenBank accession no.: AJ544218, AJ544219, AJ544220, AJ544221), we carried out a selective PCR, which amplified the HS1,2-A region, but not the identical inverted HS1,2-B region (Giambra *et al.*, 2005). Genomic DNA was extracted from peripheral blood nucleated cells and amplified with the primers described previously (Giambra *et al.*,

2005). The cycle conditions were changed to 94 °C at 2 minutes for the first step, followed by 94 °C at 30 seconds, 61 °C at 30 seconds, 68 °C for 5 minutes for 10 cycles, and 94 °C for 30 seconds, 59 °C for 30 seconds, 68 °C for 5 minutes for 20 cycles, and ending with 72 °C for 10 minutes. PCRs were carried out in 50 μ l of reaction volume containing 2 μ l of extracted DNA (50 ng), 1.5 U Platinum *Taq* DNA polymerase High Fidelity (Invitrogen, Carlsbad, CA), 15 pmol of each primer, 1.5 mM MgSO₄, 50 μ M each dNTP, and 1 \times buffer High Fidelity (600 mM Tris-SO₄, pH 8.9, 180 mM [(NH₄)₂SO₄]; Invitrogen) by using the GeneAmp PCR System 9700 (Applied Biosystems, Foster City, CA).

To prevent carryover contamination, pre-PCR procedures were performed with dedicated equipment in a laminar flow hood, using aerosol-resistant plugged pipette tips (ART; Molecular Bio-Product, San Diego, CA). Permanent devices were sterilized by ultraviolet irradiation between usages. Negative and positive controls, without DNA template or with a control DNA of a heterozygote, were always included.

A nested second PCR to amplify the polymorphic core of the enhancer HS1,2-A was performed with 1/25 of the volume of the first PCR, avoiding carryover of the genomic DNA of the first reaction. Control reactions were performed with 4 and 10 ng of total genomic DNA and resulted in no visible amplification in those conditions on agarose gel electrophoresis. The primers for the PCRs are reported by Giambra *et al.* (2005). Cycle conditions were as follows: denaturation step, 94 °C for 5 minutes; then 94 °C for 30 seconds, 56 °C for 30 seconds, 68 °C for 60 seconds, for 30 cycles; then the last cycle at 72 °C for 2 minutes (final extension step). This second PCR was carried out with the same volumes and concentrations as in the first PCR, except for the use of 1 U of Platinum *Taq* DNA polymerase (Invitrogen). PCR products were analyzed on 2.5 % agarose gel stained with ethidium bromide.

Statistical analysis

The frequencies of the variables considered are described in percentages. Comparisons between the frequencies of categorical variables in different groups were performed by χ^2 -test. *P*-value was set at 0.05. Smith's Statistical Package, version 2.80 (Pomona College, Claremont, CA), Graphpad Instat (San Diego, CA), and Stata 6.0 TM software (College Station, TX) were used for statistical analysis.

CONFLICT OF INTEREST

The authors state no conflict of interest.

ACKNOWLEDGMENTS

We are grateful to Professor Barbara K. Birshtein for correcting the previous form of the paper and important suggestions for the discussion of the results. We thank the administrative staff at the Department of Biology "E. Caffè", University of Tor Vergata, for managing funds and Dr Francesco Raducci for excellent laboratory support. This research was supported by funds from the annual grant for research from the University of Tor Vergata and the Catholic University of Roma.

REFERENCES

- Andersson T, Samuelsson A, Matthias P, Pettersson S (2000) The lymphoid-specific cofactor OBF-1 is essential for the expression of a V(H) promoter/HS1,2 enhancer-linked transgene in late B cell development. *Mol Immunol* 37:889-99

- Aupetit C, Drouet M, Pinaud E, Denizot Y, Aldigier JC, Bridoux F *et al.* (2000) Alleles of the alpha-1 immunoglobulin gene 3' enhancer control evolution of IgA nephropathy toward renal failure. *Kidney Int* 58:966-71
- Borg AA (1997) Antibodies to cytokeratins in inflammatory arthropathies. *Semin Arthritis Rheum* 27:186-95
- Bowcock AM (2005) The genetics of psoriasis and autoimmunity. *Annu Rev Genomics Hum Genet* 6:93-122
- Caspi D, Anouk M, Golan I, Paron D, Kaufman I, Wigler I *et al.* (2006) Synovial fluid levels of anti-cyclic citrullinated peptide antibodies and IgA rheumatoid factor in rheumatoid arthritis, psoriatic arthritis, and osteoarthritis. *Arthritis Rheum* 55:53-6
- Chen C, Birshtein BK (1997) Virtually identical enhancers containing a segment of homology to murine 3'IgH-E (hs1,2) lie downstream of human Ig C alpha 1 and C alpha 2 genes. *J Immunol* 159:1310-8
- Denizot Y, Pinaud E, Aupetit C, Le Morvan C, Magnoux E, Aldigier JC *et al.* (2001) Polymorphism of the human alpha1 immunoglobulin gene 3' enhancer hs1,2 and its relation to gene expression. *Immunology* 103:35-40
- Frezza D, Giambra V, Cianci R, Fruscalzo A, Giufre M, Cammarota G *et al.* (2004) Increased frequency of the immunoglobulin enhancer HS1,2 allele *2 in coeliac disease. *Scand J Gastroenterol* 39:1083-7
- Frezza D, Giambra V, Tolusso B, De Santis M, Bosello S, Vettori S *et al.* (2007) Polymorphism of immunoglobulin (Ig) enhancer element HS1,2A: allele *2 associates with systemic sclerosis. Comparison with HLA-DR and DQ alleles frequency. *Ann Rheum Dis* 66:1210-5
- Giambra V, Fruscalzo A, Giufre' M, Martinez-Labarga C, Favaro M, Rocchi M *et al.* (2005) Evolution of human IgH3'EC duplicated structures: both enhancers HS1,2 are polymorphic with variation of transcription factor's consensus sites. *Gene* 346:105-14
- Giambra V, Martinez-Labarga C, Giufre M, Modiano D, Simpori J, Gisladdottir BK *et al.* (2006) Immunoglobulin enhancer HS1,2 polymorphism: a new powerful anthropogenetic marker. *Ann Hum Genet* 70:946-50
- Harindranath N, Mills FC, Mitchell M, Meindl A, Max EE (1998) The human elk-1 gene family: the functional gene and two processed pseudogenes embedded in the IgH locus. *Gene* 221:215-24
- Hervonen K, Karell K, Holopainen P, Collin P, Partanen J, Reunala T (2000) Concordance of dermatitis herpetiformis and celiac disease in monozygous twins. *J Invest Dermatol* 115:990-3
- Kallel Sellami M, Cherif F, Zitouni M, Laadhar L, Mokni M, Hajlaoui K *et al.* (2006) Subepidermal bullous dermatosis with IgA antibodies against 200- and 280-kDa epidermal antigens after initiation of ultraviolet B therapy for psoriasis. *Br J Dermatol* 154:193-4
- Karell K, Korponay-Szabo I, Szalai Z, Holopainen P, Mustalahti K, Collin P *et al.* (2002) Genetic dissection between coeliac disease and dermatitis herpetiformis in sib pairs. *Ann Hum Genet* 66:387-92
- Lapeyre H, Courville P, Francois A, Musette P, Le Roy F, Joly P (2006) IgA necrotic renal glomerular vasculitis in pustular psoriasis. *Ann Dermatol Venereol* 133(5 Pt 1):472-3
- Lindqvist U, Kristjansson G, Pihl-Lundin I, Hagforsen E, Michaelsson G (2006) Patients with psoriatic arthritis have an increased number of lymphocytes in the duodenal mucosa in comparison with patients with psoriasis vulgaris. *J Rheumatol* 33:924-7
- Mills FC, Harindranath N, Mitchell M, Max EE (1997) Enhancer complexes located downstream of both human immunoglobulin Calpha genes. *J Exp Med* 186:845-58
- Petersson S, Cook GP, Bruggemann M, Williams GT, Neuberger MS (1990) A second B cell-specific enhancer 3' of the immunoglobulin heavy chain locus. *Nature* 344:165-8
- Pinaud E, Aupetit C, Chauveau C, Cogne M (1997) Identification of a homolog of the C alpha 3'/hs3 enhancer and of an allelic variant of the 3'IgH/hs1.2 enhancer downstream of the human immunoglobulin alpha 1 gene. *Eur J Immunol* 27:2981-5
- Pinaud E, Khamlichi AA, Le Morvan C, Drouet M, Nalesso V, Le Bert M *et al.* (2001) Localization of the 3' IgH locus elements that effect long-distance regulation of class switch recombination. *Immunity* 15:187-99
- Sardy M, Karpati S, Merkl B, Paulsson M, Smyth N (2002) Epidermal transglutaminase (TGase 3) is the autoantigen of dermatitis herpetiformis. *J Exp Med* 195:747-57
- Shepshelovich D, Shoenfeld Y (2006) Prediction and prevention of autoimmune diseases: additional aspects of the mosaic of autoimmunity. *Lupus* 15:183-90
- Sepulveda MA, Garrett FE, Price-Whelan A, Birshtein BK (2005) Comparative analysis of human and mouse 3' Igh regulatory region identifies distinctive structural features. *Mol Immunol* 42:605-15
- Stavnezer J, Amemiya CT (2004) Evolution of isotype switching. *Semin Immunol* 16:257-75
- Wandstrat A, Wakeland E (2001) The genetics of complex autoimmune diseases: non-MHC susceptibility genes. *Nat Immunol* 2:802-9
- Woo WK, McMillan SA, Watson RG, McCluggage WG, Sloan JM, McMillan JC (2004) Coeliac disease-associated antibodies correlate with psoriasis activity. *Br J Dermatol* 151:891-4
- Zadrazil J, Tichy T, Horak P, Nikorjakova I, Zima P, Krejci K *et al.* (2006) IgA nephropathy associated with psoriasis vulgaris: a contribution to the entity of 'psoriatic nephropathy'. *J Nephrol* 19:382-6